

# CATS Decrease Surgical Site Infections



## C

# lippers

### Hair Removal:

If hair must be removed from the surgical site, clippers are the best option. **Never use a razor.**

## A

# ntibiotics

### Prophylactic Antibiotics:

Antibiotics consistent with national guidelines should be administered within 1 hour of incision time and discontinued within 24 hours, in most cases.

## T

# emperature

### Normothermia:

Perioperative maintenance of normothermia (96.8–100.4° F) leads to improved patient outcomes.

## S

# ugar

### Glucose Control:

Cardiac surgery patients should have controlled 6 a.m. serum glucose ( $\leq 200$  mg/dL) on postoperative Day 1 and Day 2.

# CATS Decrease Surgical Site Infections



## Clippers

### Hair Removal:

If hair must be removed from the surgical site, clippers are the best option. **Never use a razor.**

## Antibiotics

### Prophylactic Antibiotics:

Antibiotics consistent with national guidelines should be administered within 1 hour of incision time and discontinued within 24 hours, in most cases.

## Temperature

### Normothermia:

Perioperative maintenance of normothermia (96.8–100.4° F) leads to improved patient outcomes.

## Sugar

### Glucose Control:

Cardiac surgery patients should have controlled 6 a.m. serum glucose ( $\leq 200$  mg/dL) on postoperative Day 1 and Day 2.